

Tree Species Guide for UK Agroforestry Systems

Contents

Introduction.....	3
Who is the guide for?	3
What is agroforestry?	3
Evidence base and sources of further information	4
How were the species selected?	4
Tree Species Attributes	5
Colour scheme and confidence level	5
Main products.....	6
Fruit trees: rootstocks.....	6
Timber production	6
Impact on local soil quality	7
Nutrient and organic matter accumulation.....	7
Acidification: effects on soil pH	8
Soil erosion control	9
Shade cover impacts.....	10
Livestock fodder benefits.....	10
Risks to farming operations.....	11
Carbon sequestration	11
Native status	12
Value to wildlife	12
Other environmental impacts	13
Nutrient removal and deacidification	13
Water consumption	14
Spray-drift reduction.....	15
Susceptibility to pests and disease.....	15
Climate resilience	15
Tree Species List.....	17
Species Guide	18
References	85
Authors	88
Acknowledgements	88

Species Guide Contents

Field Maple (<i>Acer campestre</i>).....	19
Sycamore (<i>Acer pseudoplatanus</i>).....	21
Italian Alder (<i>Alnus cordata</i>)	23
Common Alder (<i>Alnus glutinosa</i>)	25
Red Alder (<i>Alnus rubra</i>).....	27
Paper-bark Birch (<i>Betula papyrifera</i>).....	29
Silver Birch (<i>Betula pendula</i>)	31
Downy Birch (<i>Betula pubescens</i>).....	33
Hornbeam (<i>Carpinus betulus</i>).....	35
Sweet Chestnut (<i>Castanea sativa</i>)	37
Hazel (<i>Corylus avellana</i>).....	39
Cider Gum (<i>Eucalyptus gunnii</i>)	41
Beech (<i>Fagus sylvatica</i>)	43
Black Walnut (<i>Juglans nigra</i>)	45
Walnut (<i>Juglans regia</i>).....	47
Apple (<i>Malus domestica</i>)	49
Scots Pine (<i>Pinus sylvestris</i>)	51
Black-poplar (<i>Populus nigra</i> ssp. <i>betulifolia</i>)	53
Hybrid Poplars (<i>Populus</i> spp.)	55
Aspen (<i>Populus tremula</i>)	57
Wild Cherry (<i>Prunus avium</i>).....	59
Plum (<i>Prunus domestica</i> ssp. <i>domestica</i>).....	61
Bird Cherry (<i>Prunus padus</i>)	63
Pear (<i>Pyrus communis</i>)	65
Sessile Oak (<i>Quercus petraea</i>).....	67
Pedunculate Oak (<i>Quercus robur</i>)	69
Red Oak (<i>Quercus rubra</i>)	71
White Willow (<i>Salix alba</i>)	73
Goat Willow (<i>Salix caprea</i>)	75
Grey Willow (<i>Salix cinerea</i>).....	77
Willow varieties for SRC (<i>Salix</i> spp.).....	79
Rowan (<i>Sorbus aucuparia</i>)	81
Small-leaved Lime (<i>Tilia cordata</i>)	83

Introduction

This tree species guide aims to provide an overview of the physical characteristics, environmental tolerances, silvicultural characteristics, and ecosystem services and disservices, for a selection of 33 species of trees and shrubs that could be planted in UK agroforestry systems.

This guide has been produced following a rapid review of existing literature and databases, together with consultation with a stakeholder group. The guide comes with some important caveats, limitations and assumptions, which are discussed in this introduction.

This guide has been developed through the Defra Nature for Climate Fund (NCF) England Tree Planting Programme (ETPP) Expanding Agroforestry project.

Who is the guide for?

The guide aims to provide a simple and accessible overview of the key attributes (whether beneficial or detrimental) of tree species in UK agroforestry systems. As such, it is suitable for farmers interested in agroforestry, and other interested parties including farm advisers, foresters, and policy makers. It is envisaged that the guide will mainly be used at an early stage in the design process, e.g., to draw up a short-list of tree species. Following this, it may be advisable for those designing an agroforestry system to seek further specialist advice appropriate to the local context.

What is agroforestry?

Agroforestry is essentially 'farming with trees' and includes both the integration of trees/shrubs into farmland, and the introduction of crops or livestock into treed habitats such as woodland and orchards (Raskin and Osborn, 2019). Agroforestry can be implemented at a field scale, by integrating trees/shrubs within fields as scattered trees or lines of trees, or at the landscape scale, through the productive use of hedgerows, shelterbelts and riparian buffers. Trees within agroforestry systems can have a wide range of functions and benefits, from protection of natural resources such as soil, to product diversification such as timber, woodfuel, fruits and nuts. In many cases, agroforestry systems are designed to deliver multiple, simultaneous benefits. This guide aims to inform decision-making as to tree species selection according to the identified needs in a local context.

Evidence base and sources of further information

An accompanying Excel database aims to provide transparency as to the evidential basis for the species guide. In the database, assessments are provided for each characteristic for each species, with more explanation than the pdf guide and accompanied by numbered references where appropriate. References were sourced from the UK or north-west Europe where possible. The database has been designed so that it can be updated as an ongoing process.

The following were key sources of information, which contain useful further information on tree species selection:

- The Silviculture of Trees used in British Forestry (Savill, 2019)
- The Agroforestry Handbook (Raskin and Osborn, 2019)
- The Woodland Trust's Tree Species Handbook (Hotchkiss and Herbert, 2022)
- The Essential Tree Selection Guide (Sjoman and Anderson, 2023)
- CABI Compendium (<https://www.cabidigitallibrary.org/journal/cabicompendium/>).

In addition, a number of freely accessible online interactive tools are available to assist with site-level decision-making for species selection. These include:

- Ecological Site Classification (<http://www.forestdss.org.uk/geoforestdss/>), which is used to assess suitability of forestry species for a specified site, and also contains projected distributions and productivity of some species under future climate scenarios.
- Climate matching tool (<https://climatematch.org.uk/>), which can inform selection of climate-resilient species.
- Ammonia reduction calculator (<https://farmtreestoair.ceh.ac.uk/ammonia-reduction-calculator>), used to guide the design of shelterbelts for ammonia mitigation.

How were the species selected?

The agroforestry tree species guide focusses on a priority list of 33 tree and shrub species. The species were selected through a combination of stakeholder engagement over species from the EWCO list, and consulting the list of species within the Agroforestry Handbook (Table 18, Briggs & Knight, 2019).

Species included in the guide do not necessarily represent a 'green list' of approved or recommended species and policies and regulations may differ between UK nations. Similarly, species omitted from the guide are not necessarily unsuitable for UK agroforestry systems. A notable omission is Ash *Fraxinus excelsior* which is not included due to Ash Dieback disease, but in the past has been a popular farmland tree (for example, in the UK silvopasture trial network set up in the 1980s). If constraints relating to Ash Dieback are resolved, Ash is likely to once again become worthy of consideration in agroforestry system design.

Tree Species Attributes

For each tree species, the first page of the species guide focuses on physical characteristics, tolerances, and silvicultural properties of the trees. The second page focuses on attributes (such as ecosystem services) where tree species choice is likely to influence the productivity, environmental impacts, and resilience of the agroforestry system.

The attributes were selected based on engagement with the stakeholder group. It is also recognised that agroforestry systems provide general benefits for which tree species choice is likely less relevant. One example is the extension of grazing season observed in agroforestry systems compared with treeless pastures (McAdam *et al.*, 2018), which is likely a general benefit of integrating trees within pasture where there is no clear evidence of a role of tree species choice.

The assessment of species attributes (such as ecosystem services) draws on direct evidence for species, in addition to inference based on the physical characteristics of species, such as maximum root depth, maximum height, and canopy spread. The approach to assessing attributes is discussed in the following sections, including attributes that were considered for inclusion in the guide, but ultimately could not be included due to a lack of species-specific evidence.

Colour scheme and confidence level

Attributes are colour-coded using a traffic-light system according to whether they provide a benefit or disbenefit, as follows:

Beneficial attributes (e.g. nutrient and organic matter accumulation)	Disbenefits (e.g. acidification)
High	High
Moderate	Moderate
Low	Low

In addition, where appropriate, for each attribute value a confidence level is displayed in the guide, as follows:

- High confidence (dark circle): direct well-replicated evidence or information from reputable sources, for the species in question.
- Moderate confidence (grey circle): evidence for the species, but less reliable, e.g. limited expert opinion or a limited number of studies / limited replication.
- Low confidence (white circle): no (or very limited) direct evidence for the species, assessment primarily inferred from other tree characteristics.

Main products

Itemised under four categories:

Tree species can contribute to the following key commercial areas:

- Food, especially fruit or nut production.
- Wood, especially high-quality timber.
- Biomass, especially for woodfuel.
- Speciality products, which are typically targeted towards a local niche market.

All tree species fulfil at least one of these categories, and in many cases more than one. The list of products focusses on those which are commonly marketed, for conciseness. However, trees often bring opportunities for numerous other speciality products, such as medicinal products, edible leaves, and dyes, depending on local market opportunities.

Fruit trees: rootstocks

Fruit trees, such as Apple, Pear, Plum and Cherry, are typically grown on rootstocks to control their vigour in addition to other benefits such as disease resistance. A summary of recommended rootstocks is provided in Table 1 below. Further guidance is readily available elsewhere¹. Similarly, many cultivars are available for these species, both traditional and modern, offering different marketable products (e.g. dessert or culinary apples), taste, visual appearance, disease resistance, and harvesting times. Careful consideration should be given to cultivar selection in agroforestry systems to ensure suitable market opportunities and to fit in with seasonal labour demands (e.g., selecting late-ripening apple varieties to avoid conflicts with arable harvest²).

Table 1. Recommended rootstocks for fruit trees in agroforestry systems (adapted from information provided by Frank P Matthews).

Fruit tree	Rootstock	Size	Notes
Apple	M25	Vigorous	Vigorous planting schemes or very poor soil
	MM111	Vigorous	Vigorous planting schemes or very poor soil. Can be temperamental, generally less preferable to M25.
	MM106	Semi-vigorous	Appropriate for most schemes
	M116	Semi-vigorous	Appropriate for most schemes, similar to MM106 but with 'wet feet' resistance
Pear	Pyrus communis	Vigorous	-
	Pyrus kirchensaller	Vigorous	A more uniform rootstock from seed than P. communis
	Pyrodwarf	Semi-vigorous	Clonally produced and less suckers than P. communis or kirchensaller. The name is deceptive.

¹ E.g. https://www.frankpmatthews.com/advice/fruit_rootstocks/, <https://www.rhs.org.uk/fruit/fruit-trees/rootstocks>

² <https://agroforestry.net.eu/afinet/whitehall-farm-an-innovative-silvoarable-orchard-system-in-the-uk>

Fruit tree	Rootstock	Size	Notes
Pear / Quince	Quince A	Semi-vigorous	Delayed compatibility issues, only appropriate in some circumstances
Cherry	Colt	Semi-vigorous	Appropriate for most schemes
	F.12.1 / avium	Vigorous	Vigorous planting schemes or very poor soil
Plum family*	Brompton	Vigorous	Vigorous planting schemes or very poor soil. Limited supply.
	St Julien A	Semi-vigorous	Appropriate for most schemes
	Wavit / Weiwa	Semi-vigorous	Appropriate for most schemes
	Myrobalan B	Vigorous	Vigorous planting schemes or very poor soil.

* Plum family includes Plums, Damsons, Gages, Peaches, Nectarines and interspecific Prunus crosses.

Timber production

For high quality hardwood and timber in agroforestry systems it is essential to choose plants that are from an improved tree breeding programme. The system's effectiveness greatly depends on both using plants with the best genetic quality and utilising the correct provenance of the tree seedlings. Additionally, it is important to plant only healthy, high-quality saplings straight from the nursery; these are saplings that have a good ratio of shoot to root mass, are free from disease or injuries and do not have forks. For some species, hybrids or 'genetically improved' varieties are available which offer improved vigour and disease resistance.

To provide an indication of timber productivity in the species guide, maximum yield classes are stated from the Ecological Site Classification as cubic metres per ha of equivalent single-species stand, per year over a typical rotation. These are maximum values achievable under optimal site conditions and management. In reality, yields are likely to be substantially lower in almost all cases, but the values aim to allow a comparison of productivity between species. Average yield classes are also stated where known.

Impact on local soil quality

The impact of tree species on soil quality is assessed according to two sub-categories: (i) nutrient accumulation, including nutrient cycling, nitrogen fixation and other soil improving characteristics, (ii) effects on soil pH. Soil erosion control was also considered for inclusion in the guide, as described below, but is not currently included.

Nutrient and organic matter accumulation

Benefit, categorised as low, moderate, or high

Trees can help to return nutrients and organic matter to the upper soil layers by accessing nutrients at deeper soil layers than crop roots, which are subsequently released back into

the topsoil e.g. via litterfall and root breakdown (Isaac and Borden, 2019; Kim and Isaac, 2022). Although nutrient cycling is a complex topic, the most relevant traits include tree rooting depth, on the premise that deeper roots can access nutrients at greater depth, and canopy height and canopy spread which increases the distribution of leaf litter (Isaac and Borden, 2019; Casals *et al.*, 2014; Pardon *et al.*, 2017; Kassa *et al.*, 2022). As such, these three traits were primarily used to assess the nutrient accumulation potential of tree species.

The nutrient content (e.g. NPK ratio) of leaf litter, and its impact on soil nutrients, also depends on tree species (e.g. PuraHong *et al.*, 2014), which is another layer of complexity beyond the remit of this rapid review.

Some trees can also increase soil nutrients through nitrogen fixation. Research of nitrogen-fixing trees is surprisingly uncommon in European agroforestry systems. One such example comprised an experimental silvopasture plot in North Wales that trialled Red Alder *Alnus rubra*, showing evidence of nitrogen fixation in the absence of fertiliser application, with equivalent pasture productivity to a Sycamore *Acer pseudoplatanus* silvopasture receiving 160 kg N/ha/year (Mmolotsi, 2004; Teklehaimanot *et al.*, 2002).

The organic matter accumulation potential of tree species is therefore broadly categorised as high, moderate, or low, based on rooting depth and canopy area, determined by tree height and canopy spread (see Table 2). Nitrogen fixation is noted separately where relevant.

Table 2. Decision matrix for predicting nutrient and organic matter accumulation of tree species.

	Small canopy area (height x width <250m ²)	Moderate canopy area (height x width <400m ²)	High canopy area (height x width >400m ²)
Deep roots	Moderate	High	High
Moderate root depth	Low	Moderate	High
Shallow roots	Low	Low	Moderate

Acidification: effects on soil pH

Disbenefit, categorised as low, moderate, or high

Tree planting on agricultural land typically increases soil acidification, i.e. decreases pH (Hagen-Thorn *et al.*, 2004; De Schrijver *et al.*, 2012; Jug *et al.*, 1999). However, this effect appears to strongly depend on tree species, because of the differing chemical composition and decomposability of leaf litter (Hagen-Thorn *et al.*, 2004). Tree species with slower litter decomposition rates and lower quantities of nutrients, such as Beech *Fagus sylvatica*, are associated with the greatest acidification effects, compared with species with rapidly decomposing litter such as Lime *Tilia* spp., in forest plantations on former agricultural land (Hagen-Thorn *et al.*, 2004; De Schrijver *et al.*, 2012).

A rapid evidence review has been undertaken to assess the effect of tree species on soil pH, albeit these are largely from forestry contexts. Tree species are classified as having high, moderate or low impacts, with 'high' species having the greatest effect on soil

acidification (decrease in pH). Species categorised as 'low' would typically still have an acidifying effect, but less pronounced than in 'high' species. The categorisation was undertaken by benchmarking against relatively well-studied species included in multiple studies (see Table 3). Some species are categorised as 'unknown' where it was not possible to benchmark their acidification effect against other species, but in some of these species there is still evidence for an acidification effect after planting.

Table 3. Overview of acidification effects of relatively well studied tree species following the rapid literature review (see accompanying database for references).

Species	Acidification effect	Comments
<i>Fagus sylvatica</i>	High	Typically associated with highest acidification effects of broadleaved species
<i>Quercus robur</i>	High	Similar, or slightly less, acidification effects as <i>F. sylvatica</i>
<i>Alnus glutinosa</i>	High	Typically similar effects to native <i>Quercus</i>
<i>Acer pseudoplatanus</i>	Moderate	Intermediate between <i>Quercus</i> and <i>Tilia</i>
<i>Tilia</i> spp.	Low	Typically lowest acidification effects of broadleaved species

Soil erosion control

Not included in the guide

Control of soil erosion is often cited as one of the primary benefits of planting trees on agricultural land (Sollen-Norrlin *et al.*, 2020; Torralba *et al.*, 2016). Soil erosion can occur through water or wind. Erosion from wind can be controlled by planting an effective windbreak, which depends more on planting density and tree height than on species choice *per se* (Böhm *et al.*, 2014).

Research on the effectiveness of different tree species to control soil erosion by water is very limited. Willow *Salix* spp. and Poplar *Populus* spp. species are commonly recommended because of their rapid growth, extensive lateral root systems, and tolerance of wet soil conditions (Stokes *et al.*, 2014). Their extensive root systems are also likely to help stabilise soils. Trees with deep root systems as well as an extensive root mass are likely to be most effective at stabilising soil (Ola *et al.*, 2015; Reubens *et al.*, 2007), while leaf litter production also helps to protect soil (Castro-Díez *et al.*, 2019). However, the situation is complicated by the possibility of soil type interacting with root architecture to determine the benefit of trees on controlling soil erosion (Vannoppen *et al.*, 2017).

Although careful consideration was given to the inclusion of this attribute in the guide, especially given the significant benefits of agroforestry systems on soil erosion control (Torralba *et al.*, 2016), at this stage soil erosion control has not been included in the guide due to a lack of evidence as to the importance of root architecture, tree growth rates, and their interaction with soil types. Further research of the effects of tree root architecture on soil erosion in different soil types is therefore needed (Stokes *et al.*, 2014).

Shade cover impacts

Benefit or disbenefit depending on system; categorised under three subcategories

Tree shade can be a desirable property in terms of livestock welfare and growth, or an undesirable property in most arable or horticultural systems in the UK. The effect of trees on spatial and temporal shading patterns is complex and depends on planting arrangement (particularly in an alley cropping system) and tree management. In the species guide, the impact of shade is reported using the following three sub-categories:

- **Size of shadow**, categorised as 'low', 'moderate' or 'high'. This is determined by the canopy volume, estimated from the maximum tree height, width, and canopy shape. Based on these variables, the area of the shadow cast by the tree is estimated, and then categorised by benchmarking against the wider species list.
- **Canopy density**, categorised as open, moderately open, moderately dense, or dense.
- **Leaf emergence**, categorised as 'early' (in April), 'late' (in May), or evergreen.

Livestock fodder benefits

Benefit, categorised as low, moderate, high, or (potentially) toxic

This category assesses the potential of each species to be used as supplementary fodder for livestock. The value of tree fodder for livestock varies according to site, season, and the part of the plant, in addition to tree and livestock species. Some nutrients, such as zinc and cobalt, strongly vary among different tree species, while others (such as selenium) appear to be more dependent on site conditions with only minor differences between species (Kendall *et al.*, 2021).

Given the importance of variables such as site and season, which vary between studies, this assessment does not include numeric values for the concentration of nutrients per species, but instead reports any nutrients or minerals shown to be present at high concentrations relative to other tree species and pasture forage. Potential disbenefits from low concentrations of particular nutrients or minerals in a species are not reported, on the assumption that any individual tree species will only be a minor supplementary component of the diet, such that any shortfalls are unlikely to be problematic in the context of a wider balanced diet. However, negative aspects are stated where the literature indicates a species has palatability or toxicity concerns.

A broad assessment of the species' benefit to livestock is evaluated as high, moderate, or low, based on evidence of nutritional value, including crude protein, condensed tannins, micronutrient levels, and palatability. For example, Italian Alder *Alnus cordata* is categorised as 'moderate' value because of its high crude protein and condensed tannin content, balanced against its low palatability. Where there is significant risk of toxicity (e.g. Wild Cherry *Prunus avium* and Bird Cherry *P. padus*), this is stated instead of the high-low nutritional value. Otherwise, a more specific or limited risk of toxicity is added as a note

following the assessment of nutritional value (e.g. for Sessile Oak *Quercus petraea*, where there is some toxicity risk in certain plant parts / livestock types).

Risks to farming operations

Free text

This is a broad category which includes any other risks to productivity and farming operations that were encountered during the review. Particular attention was given to any tendency to readily spread, for example through suckering roots or prolific production of seeds that easily germinate. Other risks include allelopathy (toxicity to other plant species), risks to agricultural production other than pests and diseases, secondary hosts for pests of agricultural crops, and flammability.

Carbon sequestration

Benefit, categorised as very low, relatively low, moderate, high, or very high

Carbon sequestration in agroforestry systems is a complex topic which depends on a range of variables such as planting density, tree management, vigour of tree growth, and site characteristics including climate (Soil Association, 2023). As such, the approach taken in this guide is to provide the maximum likely achievable carbon sequestration for each tree species using the Woodland Carbon Code³, assuming that trees are planted at 3m spacing (2m for Scots Pine). A maximum possible yield class for the UK (according to the Ecological Site Classification) was assumed for each species, to provide a consistent and comparable approach accounting for differences in productivity by species. For small non-timber species such as Apple *Malus domestica*, carbon sequestration was assumed to be 'relatively low' in the short-term, and 'very low' in the medium- and long-term.

Soil carbon sequestration was not considered in the assessment, because of the lack of species-specific evidence from agroforestry systems. Other factors such as agricultural management practices, previous land use, and soil type are likely to be more relevant. However, evidence from forestry planting indicates that deciduous broadleaved species sequester significantly more soil carbon than coniferous species, with broadleaved nitrogen-fixing species having the strongest effect (Laganière *et al.*, 2010).

The carbon sequestration values provided in the database represent tonnes of CO₂ equivalent within 1 ha of trees. As such, these figures should be multiplied by tree percentage in an agroforestry system, e.g., multiply by 0.1 for an agroforestry system comprising 10% trees for an approximate estimation. However, the values provided are not intended to be a realistic assessment of carbon sequestration in agroforestry systems (given that maximum yield classes were selected), but rather provide a comparable benchmark with which to make comparisons between species.

Three timeframes for carbon sequestration are presented in the guide: short-term (20 years), medium-term (40 years), and long-term (60 years). These refer to time after tree

³ <https://woodlandcarboncode.org.uk/>

planting, assuming no coppicing, pollarding, thinning, or felling. As such, the medium-term and long-term categories are not applicable for trees managed on a coppice rotation of less than 20 years.

For each of the three timeframes, carbon sequestration is categorised by benchmarking the sequestration values described above against the wider tree species list. This means that, for example, Field Maple *Acer campestre* delivers very low carbon sequestration relative to other tree species in the guide, but could still increase carbon sequestration relative to a non-tree scenario.

Native status

Categorised as native, long-established introduction, or recent introduction

Native status is adapted from Stace's New Flora (2019), based on the following categories and sub-categories:

- Native: a species that colonised the British Isles naturally, typically long ago, from other native areas.
- Long-established (archaeophyte): a species typically associated with human activities or suspected to be introduced by humans, and present in the British Isles since at least 1500 AD. It is often uncertain whether the species is native or introduced. Further divided into the following sub-categories:
 - Denizen: behaves like a native species but suspected to be a possible human introduction (either intentionally or accidentally).
 - Cultivated: introduced by humans as crops, now persisting in the wild.
- Recent introduction (neophyte): a non-native species that arrived in the British Isles after 1500 AD, sub-categorised as:
 - Naturalised: established in the wild and not reliant on replanting.
 - Survivor: not naturalised, but persists without spreading, usually a relic of planting.

Native status sometimes varies across the UK, and in these cases, the 'highest' status is given (e.g., a species which is native to part of the UK but introduced elsewhere is categorised as 'native' for simplicity).

Value to wildlife

Benefit, categorised as relatively negligible, low, moderate, high, or very high

The value of tree species for native wildlife is derived from an article published in British Wildlife (Alexander *et al.*, 2006). In the article, the authors estimate the value of tree species for nine different categories of species assemblages, such as 'mycorrhizal fungi', 'foliage invertebrates', 'blossom for pollen and nectar' and 'epiphyte communities', using a scoring system of 1 to 5. The values were estimated based on the opinions of a selection of established and published experts. A number of assumptions were made, including that the trees are maidens (i.e., no coppicing or pollarding) with no constraints associated with commercial operations such as inputs or other ecologically harmful management.

In the species guide, the values in the article are summarised by firstly providing an overall value of the tree species for wildlife by averaging the values (1-5) across all nine assemblage categories, which are then categorised as relatively low (average value <2.5), moderate (average value >2.5 and ≤3), and high (average value >3), with a very high category used for native *Quercus* because these were substantially higher than any other species. Secondly, the assemblage(s) for which the tree species had the highest score is stated.

For species not included in the above article, their value is inferred based on their native status and similarity to species that are included in the article. For example, Italian Alder *Alnus cordata* is predicted to have low to moderate wildlife value, based on the 'moderate' score for Common Alder *A. glutinosa*, but accounting for the introduced status of Italian versus the native Common Alder.

As for the carbon values, it should be noted that tree species' value to wildlife scores are benchmarked against the wider tree species list, not relative to non-tree scenarios. For example, planting a tree species of 'relatively low' biodiversity value in an otherwise monoculture of arable or pasture could lead to substantial and significant biodiversity benefits. The extent of these benefits are likely to depend on a range of variables including management of the tree and its understorey vegetation, tree density, habitat connectivity, and management of the adjacent crop/pasture (reviewed in Kletty *et al.*, 2023).

Other environmental impacts

Free text, limited number of tree species covered

This section includes wider environmental impacts which were considered for inclusion as separate categories in the guide, but for which limited species-specific evidence is available, as described below.

Nutrient removal and deacidification

Trees can play an important role in mitigating nutrient leaching and acidification into nearby habitats, including watercourses and terrestrial plant communities adapted to low nutrient conditions. However, tree species selection is likely to play a relatively minor role, compared with hydrogeological properties including soil type and depth and water table height (Hill, 2019), in addition to tree management (T. Nisbet pers. comm.). Indeed, a meta-analysis found no significant difference in nutrient uptake among buffer strips comprising different vegetation types, including trees versus herbaceous buffers (Mayer *et al.*, 2007).

Nevertheless, some evidence suggests that Poplars and Willows are particularly efficient at removing nitrate from water (Nisbet *et al.*, 2011; Regni *et al.*, 2021), while Poplar is also particularly effective at removing atmospheric ammonia (Tang *et al.*, 2022; CEH, n.d.). These species are also able to increase nitrate uptake when present at higher concentrations (Regni *et al.*, 2021). To at least some extent, this is likely due to the rapid

growth of these species and will therefore be strongly influenced by tree management (T. Nisbet pers. comm.).

It has also been hypothesised that root architecture plays a role in nutrient accumulation, with deeper-rooting species such as Walnut *Juglans* spp. potentially more effective at reducing nitrogen leaching because of a greater 'safety net' effect (Andrianarisoa *et al.*, 2016). This hypothesis warrants further research, but there is insufficient evidence at present to include in the guide.

In some cases, trees can worsen leaching of nutrients into watercourses, and contribute to surface water acidification (Nisbet and Evans, 2014). This includes nitrogen-fixing Alder *Alnus* species, where nitrogen levels are already high, and species which consume high quantities of water coupled with limited nutrient uptake, such as Scots Pine *Pinus sylvestris*.

In terms of atmospheric nutrient removal, an online ammonia reduction tool provides advice on the design of shelterbelts for ammonia mitigation, including species selection (CEH, n.d.). Although the effect of species on ammonia recapture depends on location and soil type, Aspen *Populus tremula* is generally the best-performing species of those included in the tool. However, other factors such as buffer dimensions are more important than species selection.

Given the apparently limited role of species selection for nutrient removal and deacidification, except in some cases, no specific category is included in the species guide, and key species (whether beneficial or detrimental) such as Alder, Poplars and Willows are referred to within the 'other environmental impacts' category.

Water consumption

High water use by trees can either be beneficial in areas of excess water, such as those which are prone to flooding, or detrimental where water is in short supply and there is a risk of drought. Generally, trees typically use more water than herbaceous vegetation such as pasture or arable crops. Comparable species-specific information on water use is scarce, and plays a secondary role to climatic and soil factors, in addition to tree management (given that growing trees generally use more water than mature trees) (Nisbet, 2005; T. Nisbet pers. comm.). Therefore, water consumption by species is not included as a separate category in this guide.

Broadly speaking however, evergreen coniferous species have higher water consumption than broadleaved species, and of the broadleaved species, high-growth trees managed on a short rotation coppice have the highest consumption (Nisbet, 2005; T. Nisbet pers. comm.). In particular, Willow and Poplar have very high transpiration rates in wet soil conditions (Nisbet, 2005), but less so in drier soils (Hall *et al.*, 1996; T. Nisbet pers. comm.). Similarly, there is evidence of high water use of Common Alder *Alnus glutinosa* when soil water content is high (Herbst *et al.*, 1999). As such, the potential for high water consumption is noted for these species under 'other environmental impacts'.

Spray-drift reduction

Not included in the guide

Trees can also help to reduce spray-drift of agrochemicals onto nearby habitats. Traits such as timing of leaf emergence (Wenneker and Van de Zande, 2008) and leaf structure, hairiness and coarseness (Ucar and Hall, 2001; Ucar *et al.*, 2003; Bentrup *et al.*, 2019) appear to be important in determining the effect of a species, although other characteristics such as the height, width and porosity of a tree row or hedgerow are possibly more important (reviewed in Bentrup *et al.*, 2019; Ucar and Hall, 2001). As such, given the limited evidence at present, this category is not included in the guide.

Susceptibility to pests and disease

Disbenefit, categorised under three subcategories as low, moderate, or high

Three categories are assessed, namely susceptibility to (i) disease, (ii) invertebrate pests (e.g. insects), and (iii) vertebrate pests (e.g. mammals and birds).

For each category, a broad classification is made as follows:

- High susceptibility: the species is commonly affected by a pest or disease which causes serious damage (e.g., major loss of crop or tree mortality).
- Moderate susceptibility: the species is commonly affected by a pest or disease which causes less serious damage, i.e. trees can often make a full recovery, or less commonly affected by a serious pest/disease.
- Low susceptibility: major pest or disease issues are rare.

The above categories are generalised for the UK at the time of writing. In reality, pest and disease issues are complex and depend on numerous factors such as climate, the presence of the pest or disease in the local area either currently or in the recent past, the composition of habitats and plant species locally and in the wider landscape, and tree stress (e.g., caused by unfavourable environmental conditions). In addition, populations of tree pests and diseases are rapidly changing and there is a constant threat of new pests and diseases colonising the UK, primarily due to climate change and global trade. It is advisable that a mixture of species, or at least varieties, are planted in agroforestry systems to help mitigate this threat.

Climate resilience

Benefit, categorised as low, moderate or high

The resilience of each tree species to projected climate change was assessed by undertaking a rapid review of the literature for each species, e.g. using the search terms “*Juglans regia* climate change UK”, in addition to reviewing grey literature. The identified studies typically assess climate resilience by modelling species’ environmental tolerances such as temperature requirements and drought tolerance against projected 2050 climate change scenarios, and do not typically account for any changes in community dynamics, for example from pest pressure or competition from other tree species. Where

species-specific evidence was not found in the literature, a prediction is made based on the environmental tolerances of the species and its native distribution. The resilience of each species is evaluated according to three categories based on the available evidence:

- High: UK climate likely to generally become more suitable for the species, which is projected to expand its range within the UK with very limited areas of reduced suitability.
- Moderate: mixed effects of projected climate change in the UK, e.g. projected expansion in the north, but with reduced suitability in a comparable area in the south.
- Low: projected climate change likely to result in substantially reduced suitability for the species in the UK, with limited range expansion.

For example, Sycamore *Acer pseudoplatanus* is categorised as 'moderate' because the species is likely to become less suited to the south-east, east and midlands of England, but more suited to northern England and south-east Scotland, such that its overall range would cover a similar area.

Note that each species is broadly assessed across the UK. In reality, the effect of climate change on species' suitability is often strongly dependent on the region of the UK.

Tree Species List

Common Name	Latin Name
Field Maple	<i>Acer campestre</i>
Sycamore	<i>Acer pseudoplatanus</i>
Italian Alder	<i>Alnus cordata</i>
Common Alder	<i>Alnus glutinosa</i>
Red Alder	<i>Alnus rubra</i>
Paper-bark Birch	<i>Betula papyrifera</i>
Silver Birch	<i>Betula pendula</i>
Downy Birch	<i>Betula pubescens</i>
Hornbeam	<i>Carpinus betulus</i>
Sweet Chestnut	<i>Castanea sativa</i>
Hazel	<i>Corylus avellana</i>
Cider Gum	<i>Eucalyptus gunnii</i>
Beech	<i>Fagus sylvatica</i>
Black Walnut	<i>Juglans nigra</i>
Walnut	<i>Juglans regia</i>
Apple	<i>Malus domestica</i> / cultivars
Scots Pine	<i>Pinus sylvestris</i>
Black-poplar	<i>Populus nigra</i> ssp. <i>betulifolia</i>
Hybrid poplars (timber)	<i>Populus</i> spp.
Aspen	<i>Populus tremula</i>
Wild Cherry	<i>Prunus avium</i>
Plum	<i>Prunus domestica</i> ssp. <i>domestica</i>
Bird Cherry	<i>Prunus padus</i>
Pear	<i>Pyrus communis</i>
Sessile Oak	<i>Quercus petraea</i>
Pedunculate Oak	<i>Quercus robur</i>
Red Oak	<i>Quercus rubra</i>
White Willow	<i>Salix alba</i>
Goat Willow	<i>Salix caprea</i>
Grey Willow	<i>Salix cinerea</i>
Willow varieties (SRC)	<i>Salix</i> spp.
Rowan	<i>Sorbus aucuparia</i>
Small-leaved Lime	<i>Tilia cordata</i>

Species Guide

Field Maple (*Acer campestre*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Typically up to 15m , exceptionally 25m
Canopy cover	Rounded 4-8m wide
Canopy density	Dense
Root architecture	Very shallow
Growth rate	Moderate to high
Silviculture	
Establishment time	10-20 years to maximum height
Establishment requirements	Protection from browsing
Management requirements	Minimal to no pruning
Longevity	Usually up to 120 years , potentially 200+ years
Rotation length	Typically 8-15 years , up to 30 years
Approach to silviculture	Coppices well
Tolerances	
High temperatures	Tolerant
Low temperatures	Hardy
Wind	Tolerant
Drought	Moderately tolerant
Waterlogging	Moderately sensitive to sensitive
Soil types	Neutral to calcareous, moist but well-drained
Slope and aspect	Any
Shade tolerance	Full sun or partial shade

Productivity		
Main products	Food	-
	Wood	Timber, low yielding
	Biomass	Woodfuel
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Moderate ○
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Low to moderate (low crude protein, moderate nutritional value) ●
Risks to farming operations		Could inhibit growth of nearby plants
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Native ●
Value to wildlife		Relatively low, highest value for pollen & nectar ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Low to moderate ●
	Invertebrates	Moderate ●
	Vertebrates	High, especially Rabbits ●
Climate resilience		High ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Sycamore (*Acer pseudoplatanus*)

Physical	
Typical systems	Arable Pasture Lowland and upland
Max. height	Typically up to 35m , exceptionally 40m
Canopy cover	Broad, domed to ovoid >8m wide
Canopy density	Moderate when young, otherwise dense
Root architecture	Shallow to moderate, extensive
Growth rate	Moderate to high
Silviculture	
Establishment time	> 50 years to maximum height
Establishment requirements	Protection from browsing Benefits from weed control
Management requirements	Minimal to no pruning, self-prunes well
Longevity	400 or 500 years
Rotation length	Typically 65-75 year timber rotation, up to 100 years
Approach to silviculture	Can be coppiced up to 80-100 years Can be pollarded
Tolerances	
High temperatures	Moderately sensitive
Low temperatures	Very hardy
Wind	Tolerant, but benefits from shelter
Drought	Moderately sensitive
Waterlogging	Moderately sensitive to sensitive
Soil types	Wide range, preferring deep well-drained soils
Slope and aspect	Any
Shade tolerance	Full sun or partial shade

Productivity		
Main products	Food	-
	Wood	Timber, moderately high yielding
	Biomass	Woodfuel
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	Moderate ●
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		Low to moderate (conflicting evidence for crude protein, low digestibility) ● Risk of toxicity in horses
Risks to farming operations		Self-sows freely, potential weed
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Recent introduction (Neophyte – naturalised) ●
Value to wildlife		High, especially foliage invertebrates, leaf litter, epiphytes ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Moderate to high ●
	Invertebrates	Moderate ●
	Vertebrates	High (especially squirrels) ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Italian Alder (*Alnus cordata*)

Physical	
Typical systems	Arable Lowland
Max. height	20 to 29m
Canopy cover	Conical 4-8m wide
Canopy density	Moderately dense
Root architecture	Shallow
Growth rate	High
Silviculture	
Establishment time	20 to 50 years to maximum height
Establishment requirements	Minimal
Management requirements	Minimal to no pruning
Longevity	60 to 100 years
Rotation length	At least 20-30 year timber rotation
Approach to silviculture	Variable coppicing ability, rotation 15-30 years
Tolerances	
High temperatures	Moderately tolerant
Low temperatures	Hardy
Wind	Tolerant
Drought	Tolerant, but reduces growth in drought
Waterlogging	Moderately tolerant
Soil types	Prefers deep chalky soils
Slope and aspect	Any
Shade tolerance	Full sun or partial shade

Productivity		
Main products	Food	-
	Wood	Timber, pulpwood, moderately high yielding
	Biomass	Firewood
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Nitrogen fixing ● Otherwise low ○
	Acidification	High ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Moderately dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		Moderate (high protein content, low palatability) ●
Risks to farming operations		None known
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Recent introduction (Neophyte – naturalised) ●
Value to wildlife		Low to moderate ○
Other environmental impacts		Risk of increased nitrate leaching and acidification
Resilience		
Pest/disease susceptibility	Diseases	Moderate ●
	Invertebrates	Low ●
	Vertebrates	Low ○
Climate resilience		High ○

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Common Alder (*Alnus glutinosa*)

Physical	
Typical systems	Arable Pasture Lowland to semi-upland
Max. height	Typically 12 to 25m , exceptionally 40m
Canopy cover	Broad, conical to ovoid 4-10m wide
Canopy density	Open
Root architecture	Moderate, to very shallow in wet soils
Growth rate	Moderate to high
Silviculture	
Establishment time	20 to 50 years to maximum height
Establishment requirements	Minimal
Management requirements	Minimal to no pruning
Longevity	Typically 100 years, potentially >250 years, only 20-25 years on poor sites
Rotation length	Maximum rotation 60-70 years
Approach to silviculture	Coppices well when young Can be pollarded
Tolerances	
High temperatures	Moderately tolerant
Low temperatures	Very hardy
Wind	Tolerant, but shelter required for straight stems
Drought	Moderately sensitive to sensitive, especially when young
Waterlogging	Tolerant
Soil types	Requires moist soils, otherwise undemanding
Slope and aspect	Any
Shade tolerance	Full sun or partial shade

Productivity		
Main products	Food	-
	Wood	Low quality timber, moderately high yielding
	Biomass	Biomass, charcoal
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Nitrogen fixing ● Otherwise low ○
	Acidification	High ●
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate (high crude protein, moderate digestibility, low palatability) ●
Risks to farming operations		Potential weed, readily self-sows
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Native ●
Value to wildlife		Moderate, highest value for foliage invertebrates, fruits & seeds ●
Other environmental impacts		Risk of increased nitrate leaching and acidification High water consumption in wet conditions
Resilience		
Pest/disease susceptibility	Diseases	Moderate ●
	Invertebrates	Low ●
	Vertebrates	Low to moderate ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Red Alder (*Alnus rubra*)

Physical	
Typical systems	Pasture Lowland
Max. height	Typically to 15m , exceptionally 25m
Canopy cover	Narrow, pyramidal 4-8m wide
Canopy density	Moderately open
Root architecture	Moderate, extensive
Growth rate	High
Silviculture	
Establishment time	20 to 50 years to maximum height
Establishment requirements	Minimal
Management requirements	Minimal to no pruning
Longevity	Up to 60-100 years, but typically begins to die back after 15 years
Rotation length	Pulpwood 10-12 years, sawlog 30-32 years
Approach to silviculture	Coppices well on short cycles when young
Tolerances	
High temperatures	Sensitive
Low temperatures	Very hardy, but susceptible to spring frosts
Wind	Tolerant
Drought	Tolerant when established
Waterlogging	Unknown
Soil types	Moist but well-drained
Slope and aspect	Avoid north-facing
Shade tolerance	Full sun

Productivity		
Main products	Food	-
	Wood	Timber and pulpwood, moderately high yielding
	Biomass	-
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Nitrogen fixing ● Otherwise low ○
	Acidification	High ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Moderately open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Poorly understood, likely low palatability ○
Risks to farming operations		Poor growth in most UK trials
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Recent introduction (Neophyte – naturalised) ●
Value to wildlife		Low to moderate ○
Other environmental impacts		Risk of increased nitrate leaching and acidification
Resilience		
Pest/disease susceptibility	Diseases	Moderate ●
	Invertebrates	Low ●
	Vertebrates	Low ○
Climate resilience		High ○

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Paper-bark Birch (*Betula papyrifera*)

Physical	
Typical systems	Lowland
Max. height	Typically to 23-25m , exceptionally 30m
Canopy cover	Ovoid, spreading >8m wide
Canopy density	Open
Root architecture	Shallow
Growth rate	High
Silviculture	
Establishment time	20 to 50 years to maximum height
Establishment requirements	Regular mulching / weed control
Management requirements	Minimal to no pruning
Longevity	Up to 200 years
Rotation length	Likely 30-40 year timber rotation
Approach to silviculture	Coppices weakly Pollards weakly
Tolerances	
High temperatures	Sensitive
Low temperatures	Very hardy
Wind	Tolerant
Drought	Sensitive
Waterlogging	Sensitive
Soil types	Wide range, favours deep, fertile and well aerated soils
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	-
	Wood	Pulpwood
	Biomass	-
	Speciality	Tree sap
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Conflicting evidence
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Poorly understood ○
Risks to farming operations		Highly flammable bark
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Moderate
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Relatively low
Native status		Recent introduction (Neophyte – survivor) ●
Value to wildlife		Moderate to high ○
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Low (high future risk) ●
	Vertebrates	Low to moderate ○
Climate resilience		High ○

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Silver Birch (*Betula pendula*)

Physical	
Typical systems	Pasture Arable Lowland to semi-upland
Max. height	Typically to 15-20m , exceptionally 30m
Canopy cover	Columnar, tapering 4-8+m wide
Canopy density	Open
Root architecture	Shallow, deeper on dry sites
Growth rate	High
Silviculture	
Establishment time	20 to 50 years to maximum height
Establishment requirements	Regular mulching / weed control Protection from browsing
Management requirements	Minimal to no pruning
Longevity	Typically up to 100 years , exceptionally 200 years
Rotation length	30-40 year timber rotation
Approach to silviculture	Coppices weakly
Tolerances	
High temperatures	Sensitive
Low temperatures	Very hardy
Wind	Tolerant
Drought	Sensitive
Waterlogging	Sensitive
Soil types	Favours light, well-drained, acid soils
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	-
	Wood	Usually pulpwood, moderate to low yielding
	Biomass	-
	Speciality	Tree sap
Impact on local soil quality	Nutrient and organic matter accumulation	Generally low ○ Useful for restoration of acid soils ●
	Acidification	Conflicting evidence
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Low to moderate ○
Risks to farming operations		None known
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Native ●
Value to wildlife		High, especially mycorrhizal fungi and foliage invertebrates ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Low (high future risk) ●
	Vertebrates	Low to moderate ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Downy Birch (*Betula pubescens*)

Physical	
Typical systems	Pasture Lowland and upland
Max. height	Typically up to 20m , exceptionally 30m
Canopy cover	Irregular 6-8m wide
Canopy density	Open
Root architecture	Shallow to moderate
Growth rate	Moderate to high
Silviculture	
Establishment time	20 to 50 years to maximum height
Establishment requirements	Regular mulching / weed control Protection from browsing
Management requirements	Minimal to no pruning
Longevity	Typically up to 100 years , exceptionally 200 years
Rotation length	30-40 year timber rotation
Approach to silviculture	Coppices weakly, long rotation
Tolerances	
High temperatures	Sensitive
Low temperatures	Very hardy
Wind	Tolerant
Drought	Sensitive
Waterlogging	Moderately tolerant
Soil types	Wide range, favours more acid, wet, peaty soils
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	-
	Wood	Timber and pulpwood, relatively low yielding
	Biomass	Woodfuel
	Speciality	Tree sap
Impact on local soil quality	Nutrient and organic matter accumulation	Generally low ○ Useful for restoration of acid soils ●
	Acidification	Conflicting evidence
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Low to moderate ○
Risks to farming operations		None known
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Moderate
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Relatively low
Native status		Native ●
Value to wildlife		High , especially mycorrhizal fungi and foliage invertebrates ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Low (high future risk) ●
	Vertebrates	Low to moderate ●
Climate resilience		Low ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Hornbeam (*Carpinus betulus*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Typically up to 25m , exceptionally 32m
Canopy cover	Ovoid to globular >8m wide
Canopy density	Dense
Root architecture	Shallow to moderate
Growth rate	Low to moderate
Silviculture	
Establishment time	> 50 years to maximum height
Establishment requirements	Protection from squirrel damage
Management requirements	Minimal to no pruning
Longevity	Potentially 400+ years
Rotation length	15-25 year to 30-40 year coppice rotation
Approach to silviculture	Coppices well Can be pollarded
Tolerances	
High temperatures	Tolerant
Low temperatures	Very hardy
Wind	Tolerant
Drought	Moderately tolerant
Waterlogging	Sensitive
Soil types	Wide range, favours moderately fertile, damp soils
Slope and aspect	Any
Shade tolerance	Full sun to at least partial shade

Productivity		
Main products	Food	-
	Wood	Specialised timber, moderate yielding
	Biomass	Woodfuel, charcoal
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low to moderate ○
	Acidification	High ●
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Low (moderate crude protein, low nutritional value) ●
Risks to farming operations		None known
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Moderate
Native status		Native ●
Value to wildlife		Relatively low, highest value for mycorrhizal fungi, leaf litter, seeds ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Low ●
	Vertebrates	Moderate to high (especially squirrels) ●
Climate resilience		High ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Sweet Chestnut (*Castanea sativa*)

Physical	
Typical systems	Arable Pasture Lowland to semi-upland
Max. height	30 to 35m
Canopy cover	Ovoid to irregular >8m wide
Canopy density	Dense
Root architecture	Moderate to deep, extensive
Growth rate	Moderate to high
Silviculture	
Establishment time	20-50 years to maximum height, full nut production in 10-15 years
Establishment requirements	Minimal
Management requirements	Minimal to no pruning
Longevity	Potentially 300-1000 years
Rotation length	12-16 years coppice, 25-30 years pulpwood
Approach to silviculture	Coppices well
Tolerances	
High temperatures	Tolerant
Low temperatures	Hardy (but cultivars can be sensitive to frost)
Wind	Potentially sensitive
Drought	Moderately sensitive to tolerant (cultivars typically more sensitive)
Waterlogging	Sensitive
Soil types	Wide range, favours acid to neutral, moist but well-drained soils
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	Nuts, typically using hybrid cultivars
	Wood	Timber, pulpwood, poles, moderately high yielding
	Biomass	Woodfuel (low quality)
	Speciality	Tannins for leather etc., mushroom cultivation
Impact on local soil quality	Nutrient and organic matter accumulation	High ○ Useful soil improver on light soils ●
	Acidification	Unknown
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		Conflicting evidence. Nuts used for pig feed.
Risks to farming operations		Timber often affected by ring shake
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Moderate
Native status		Long-established (Archaeophyte – cultivated) ●
Value to wildlife		Relatively low, highest value for seeds ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Low ●
	Vertebrates	Moderate ●
Climate resilience		Moderate to high ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Hazel (*Corylus avellana*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Typically up to 6m , potentially 10m
Canopy cover	Globular, spreading 4-8m wide
Canopy density	Moderately dense
Root architecture	Shallow, limited extent
Growth rate	Moderate
Silviculture	
Establishment time	5-10 years to maximum height, full nut production from c. 10 years
Establishment requirements	Protection from browsing
Management requirements	Minimal to no pruning
Longevity	Typically 70-80 years , potentially 200+ years
Rotation length	6-9 years coppice, or 14-16 years with standard trees
Approach to silviculture	Coppices well
Tolerances	
High temperatures	Moderately tolerant
Low temperatures	Hardy
Wind	Tolerant
Drought	Moderately sensitive
Waterlogging	Sensitive
Soil types	Wide range, favours base-rich, damp but well-drained soils
Slope and aspect	Avoid north facing
Shade tolerance	Favours full sun, but shade tolerant

Productivity		
Main products	Food	Nuts
	Wood	Fencing poles
	Biomass	Woodfuel
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Unknown
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Moderately dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Low (low crude protein, low digestibility) ●
Risks to farming operations		Suckering roots (rootstocks can mitigate this)
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Native ●
Value to wildlife		Moderate, highest value leaf litter and epiphytes ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Low ●
	Invertebrates	Low ●
	Vertebrates	Moderate to high (especially squirrels) ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Cider Gum (*Eucalyptus gunnii*)

Physical	
Typical systems	Lowland
Max. height	25 to 34m
Canopy cover	Ovoid >8m wide
Canopy density	Open
Root architecture	Poorly understood, potentially extensive
Growth rate	High to very high
Silviculture	
Establishment time	10-20 years to maximum height
Establishment requirements	Weed control
Management requirements	Minimal to no pruning
Longevity	Unknown
Rotation length	Typically 12 year rotation
Approach to silviculture	Can be coppiced Can be pollarded
Tolerances	
High temperatures	Sensitive
Low temperatures	Hardy in most of UK, but susceptible to early frosts
Wind	Sensitive
Drought	Likely moderately sensitive
Waterlogging	Moderately sensitive to moderately tolerant
Soil types	Favours slightly acidic
Slope and aspect	Avoid exposed locations
Shade tolerance	Full sun

Productivity		
Main products	Food	-
	Wood	Timber, pulpwood, potentially very high yielding
	Biomass	Woodfuel
	Speciality	Oil
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	Unknown
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Open ●
	Leaf emergence	Evergreen ●
Livestock fodder benefits		Limited evidence
Risks to farming operations		High risk of failure from wind-chill and frosts Highly flammable
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Very high
	Medium-term (40 yrs)	Very high
	Long-term (60 yrs)	Very high
Native status		Recent introduction (Neophyte – survivor) ●
Value to wildlife		Negligible, but some value for bees and other pollinators ●
Other environmental impacts		High water consumption
Resilience		
Pest/disease susceptibility	Diseases	Moderate ●
	Invertebrates	Low ●
	Vertebrates	Low ●
Climate resilience		Moderate to high ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Beech (*Fagus sylvatica*)

Physical	
Typical systems	Pasture Arable Lowland to upland
Max. height	Typically up to 35m , exceptionally 43m
Canopy cover	Globular to broad ovoid >8m wide
Canopy density	Dense
Root architecture	Shallow to moderate, extensive
Growth rate	Moderate
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Weed control
Management requirements	Minimal to no pruning
Longevity	Typically up to 150-300 years , exceptionally 900 years
Rotation length	Typically 70-80 year timber rotation, exceptionally 100-140 years
Approach to silviculture	Coppices weakly Can be pollarded
Tolerances	
High temperatures	Sensitive
Low temperatures	Hardy, but susceptible to early and late frosts
Wind	Tolerant, but benefits from shelter when young
Drought	Moderately sensitive
Waterlogging	Sensitive
Soil types	Wide range, but requires moderate moisture levels
Slope and aspect	Any
Shade tolerance	Full sun to at least partial shade

Productivity		
Main products	Food	-
	Wood	Timber, moderate yielding
	Biomass	Woodfuel
	Speciality	Oil from nuts
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	High ●
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Low to moderate (moderate crude protein and micronutrients) ●
Risks to farming operations		Potentially very large tree
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Moderate
Native status		Native ●
Value to wildlife		High, especially for mycorrhizal fungi, wood-decay fungi and invertebrates, seeds, epiphytes ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Moderate ●
	Invertebrates	Moderate ●
	Vertebrates	High (especially squirrels) ●
Climate resilience		Low to moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Black Walnut (*Juglans nigra*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Typically up to 30m , exceptionally 50m
Canopy cover	Globular, domed >8m wide
Canopy density	Moderate
Root architecture	Deep
Growth rate	Moderate
Silviculture	
Establishment time	> 50 years to maximum height, 10 years for commercial nut yields
Establishment requirements	Formative pruning Weed control
Management requirements	High pruning for timber
Longevity	Unknown
Rotation length	Typically 60 year timber rotation
Approach to silviculture	Can be coppiced Can be pollarded
Tolerances	
High temperatures	Tolerant
Low temperatures	Hardy, but susceptible to late frosts
Wind	Highly sensitive
Drought	Moderately sensitive
Waterlogging	Sensitive
Soil types	Somewhat demanding, favours deep, well-drained, fertile soils
Slope and aspect	Favours south / south-west, sheltered locations. Avoid north facing.
Shade tolerance	Full sun

Productivity		
Main products	Food	Nuts, more difficult to de-shell than <i>J. regia</i>
	Wood	High value timber, moderately high yielding. Vigorous hybrids available (likely lower value)
	Biomass	-
	Speciality	Dye, oil
Impact on local soil quality	Nutrient and organic matter accumulation	High ○
	Acidification	Low ○
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Moderate ●
	Leaf emergence	Late ●
Livestock fodder benefits		Moderate to high (high crude protein, moderate digestibility) ● Risk of toxicity to horses from wood shavings and leaf ingestion, and all animals from mouldy walnuts
Risks to farming operations		Allelopathic toxicity to many other plants Limited climatic suitability at present
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Moderate
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Relatively low
Native status		Recent introduction (Neophyte – naturalised) ●
Value to wildlife		Relatively low, greatest value for mycorrhizal fungi and leaf litter ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Moderate to high ●
	Invertebrates	Moderate ●
	Vertebrates	High (especially squirrels) ●
Climate resilience		High ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Walnut (*Juglans regia*)

Physical				
Typical systems	Arable	Pasture	Lowland	
Max. height	20-30m			
Canopy cover	Globular to ovoid or domed (conical when young)		Up to 15m wide	
Canopy density	Moderate			
Root architecture	Very deep			
Growth rate	Moderate			
Silviculture				
Establishment time	20-50 years to maximum height, nut yields after 5 to 6 years			
Establishment requirements	Staking	Irrigation	Formative pruning for timber	Weed control
Management requirements	High pruning for timber			
Longevity / rotation length	150-280 years			
Rotation length	Typically 60 years, exceptionally 30 years			
Approach to silviculture	Can be pollarded			
Tolerances				
High temperatures	Tolerant			
Low temperatures	Hardy, but sensitive to unseasonable frosts (some varieties have improved frost resistance)			
Wind	Sensitive (flowers and foliage)			
Drought	Moderately sensitive			
Waterlogging	Sensitive			
Soil types	Demanding; favours well-drained, deep, fertile, alkaline loam			
Slope and aspect	Avoid exposed locations			
Shade tolerance	Full sun. Light shade tolerated in early development			

Productivity		
Main products	Food	Nuts (typically using specific varieties)
	Wood	High-value timber or sawnwood, relatively low yielding. Vigorous hybrids available (likely lower value).
	Biomass	-
	Speciality	Oil, dye
Impact on local soil quality	Nutrient and organic matter accumulation	High ○
	Acidification	Low ○
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Moderate ●
	Leaf emergence	Late ●
Livestock fodder benefits		Moderate to high (high protein content, moderate digestibility) ● Risk of toxicity to horses from wood shavings, and all animals from mouldy walnuts
Risks to farming operations		Allelopathic toxicity to other plants, particularly affecting tomatoes and apples Limited climatic suitability at present
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Moderate
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Relatively low
Native status		Long-established (Archaeophyte – cultivated) ●
Value to wildlife		Relatively low, highest value for mycorrhizal fungi and leaf litter ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Low ●
	Invertebrates	Low, although nuts susceptible to pest damage ●
	Vertebrates	High (especially squirrels) ●
Climate resilience		High ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Apple (*Malus domestica*)

Physical				
Typical systems	Arable	Pasture	Lowland	
Max. height	Depends on cultivar and rootstock			
Canopy cover	Ovoid to irregular crown	Spread depends on cultivar / rootstock		
Canopy density	Moderate			
Root architecture	Depends on cultivar and rootstock, generally very deep for its size			
Growth rate	Generally moderate (depends on cultivar and rootstock)			
Silviculture				
Establishment time	Full cropping in 3-9 years (depending on cultivar and rootstock)			
Establishment requirements	Staking	Formative pruning	Water during drought	Protection from browsing
Management requirements	Annual pruning			
Longevity	Depends on cultivar and rootstock			
Rotation length	12-45 years , depending on cultivar and rootstock			
Approach to silviculture	Orchard tree			
Tolerances				
High temperatures	Tolerant			
Low temperatures	Sensitive to frost pockets			
Wind	Sensitive			
Drought	Moderately sensitive			
Waterlogging	Sensitive			
Soil types	Wide range, avoid poorly drained or shallow soils			
Slope and aspect	Sunny and sheltered			
Shade tolerance	Full sun			

Productivity		
Main products	Food	Fruits (inc. eating, cooking, cider)
	Wood	Speciality timber
	Biomass	Limited woodfuel from pruning
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	Unknown
Shade cover impacts*	Size of shadow (full-grown)	Low, but depends on rootstock/variety ●
	Canopy density	Moderate ●
	Leaf emergence	Early ●
Livestock fodder benefits		High (especially fruits and pomace) ●
Risks to farming operations		None known
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Long-established (Archaeophyte – denizen or cultivated) ●
Value to wildlife		High, especially for foliage invertebrates, leaf litter, pollen & nectar, fruits ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Generally high (varies according to variety and rootstock) ●
	Invertebrates	Generally high (varies according to variety and rootstock) ●
	Vertebrates	Moderate ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Scots Pine (*Pinus sylvestris*)

Physical	
Typical systems	Arable Pasture Lowland to upland
Max. height	35-40m
Canopy cover	Conical, broadening with age >8m wide
Canopy density	Dense
Root architecture	Deep
Growth rate	Low
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Protection from browsing
Management requirements	Regular pruning for timber
Longevity	250-400 years
Rotation length	Typically 50-60 year timber rotation, up to 100 years
Approach to silviculture	High forest tree
Tolerances	
High temperatures	Tolerant
Low temperatures	Very hardy
Wind	Tolerant
Drought	Tolerant
Waterlogging	Moderately sensitive to moderately tolerant
Soil types	Favours well-drained, non-calcareous soils, tolerant of poor fertility
Slope and aspect	Any
Shade tolerance	Favours full sun especially when young

Productivity		
Main products	Food	Nuts
	Wood	Diverse timber uses, high yielding
	Biomass	-
	Speciality	Resin, pine oil, Christmas trees
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	High ●
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Dense ●
	Leaf emergence	Evergreen ●
Livestock fodder benefits		Moderate ●
Risks to farming operations		Capable of becoming very large Can aggressively regenerate from seeds Flammable
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Relatively low
	Long-term (60 yrs)	Moderate
Native status		Native (in Scotland) ●
Value to wildlife		Moderate, greatest value for mycorrhizal fungi ●
Other environmental impacts		High water consumption Can increase acidification of nearby watercourses, especially in dry areas
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Moderate ●
	Vertebrates	High ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Black-poplar (*Populus nigra* ssp. *betulifolia*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	30-40m
Canopy cover	Spreading, ovoid to irregular >8m wide
Canopy density	Moderately dense
Root architecture	Moderate to deep, vigorously suckering
Growth rate	High
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Protection from browsing
Management requirements	Regular pruning for timber Sucker removal in autumn/winter
Longevity	200-300+ years
Rotation length	Typically 6-20 years
Approach to silviculture	Can be coppiced Can be pollarded
Tolerances	
High temperatures	Moderately tolerant to tolerant
Low temperatures	Hardy
Wind	Tolerant
Drought	Sensitive to moderately sensitive
Waterlogging	Moderately tolerant to tolerant
Soil types	Favours lowland floodplains, especially loam, chalk or sand
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	-
	Wood	Low-quality timber, high yielding
	Biomass	Charcoal
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	High ○
	Acidification	Low to moderate ○
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Moderately dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Likely moderate (moderately high crude protein content) ○
Risks to farming operations		Vigorous, suckering roots Risk of overwintering aphid pests of vegetables
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Very high
	Medium-term (40 yrs)	Very high
	Long-term (60 yrs)	Very high
Native status		Native ●
Value to wildlife		Relatively low, but of high value as an endangered native tree, greatest value for foliage invertebrates ●
Other environmental impacts		High water consumption in wet conditions High potential to reduce nutrient leaching
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Moderate ●
	Vertebrates	High ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Hybrid Poplars (*Populus* spp.)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Depends on variety
Canopy cover	Depends on variety
Canopy density	Depends on variety
Root architecture	Moderate to deep, extensive surface roots
Growth rate	Very high
Silviculture	
Establishment time	Depends on variety
Establishment requirements	Weed control
Management requirements	Regular pruning
Longevity	N/A, harvested on rapid rotation
Rotation length	Typically 30-40 years , exceptionally 22 years
Approach to silviculture	High forest tree
Tolerances	
High temperatures	Tolerant
Low temperatures	Susceptible to late frosts
Wind	Moderately tolerant but benefits from shelter
Drought	Sensitive (in terms of reducing growth rate)
Waterlogging	Sensitive (in terms of reducing growth rate)
Soil types	Demanding in terms of rapid growth, requiring highly fertile, base-rich, well-drained and aerated soils
Slope and aspect	Any
Shade tolerance	Full sun

Productivity		
Main products	Food	-
	Wood	Timber, potentially very high yielding
	Biomass	Bioenergy
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	Low to moderate ○
Shade cover impacts*	Size of shadow (full-grown)	Moderate to high (depends on variety) ●
	Canopy density	Depends on variety
	Leaf emergence	Early ●
Livestock fodder benefits		Likely moderate ○
Risks to farming operations		Risk of overwintering aphid pests of vegetables
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Very high
	Medium-term (40 yrs)	Very high
	Long-term (60 yrs)	Very high
Native status		Hybrids typically derived from non-native species ●
Value to wildlife		Relatively low, greatest value for foliage invertebrates ●
Other environmental impacts		High water consumption in wet conditions High potential to reduce nutrient leaching
Resilience		
Pest/disease susceptibility	Diseases	High (but depends on variety) ●
	Invertebrates	High (but depends on variety) ●
	Vertebrates	High ●
Climate resilience		Moderate to high (depending on variety) ○

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Aspen (*Populus tremula*)

Physical	
Typical systems	Arable Pasture Lowland to upland
Max. height	18-40m
Canopy cover	Ovoid to globular, slightly conical >8m wide
Canopy density	Light to moderately dense
Root architecture	Moderate to deep, abundant suckers
Growth rate	High
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Protection from browsing
Management requirements	Minimal to no pruning, self-prunes well
Longevity	Up to 100 years
Rotation length	Coppice rotation of 20 years for pulpwood
Approach to silviculture	Coppices well within first 5 years
Tolerances	
High temperatures	Sensitive
Low temperatures	Very hardy
Wind	Tolerant
Drought	Moderately sensitive
Waterlogging	Moderately sensitive to tolerant
Soil types	Wide range, favours free-draining mineral soils
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	-
	Wood	Timber – veneers, pulp, charcoal, potentially high yielding
	Biomass	Potential for biomass where growth is rapid
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	Low to moderate ○
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Light to moderately dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		Moderate ○
Risks to farming operations		Abundant root suckers, can damage drains and buildings
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Native ●
Value to wildlife		Relatively low, greatest value for foliage invertebrates ●
Other environmental impacts		High water consumption in wet conditions High potential to reduce nutrient leaching
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Moderate ●
	Vertebrates	High ●
Climate resilience		Moderate to high ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Wild Cherry (*Prunus avium*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Typically up to 25m , exceptionally 32m
Canopy cover	Domed, globular to broad ovoid c.8m wide or more
Canopy density	Moderately dense
Root architecture	Moderate, becoming shallower with age, suckering
Growth rate	Moderate to high
Silviculture	
Establishment time	20-50 years to maximum height, fruit yields within 5-6 years
Establishment requirements	Weed control
Management requirements	High pruning for timber
Longevity	Up to 100 years or more
Rotation length	40-80 years for timber
Approach to silviculture	Coppices poorly
Tolerances	
High temperatures	Moderately sensitive
Low temperatures	Hardy, but flowers can be damaged by late frosts
Wind	Conflicting information
Drought	Moderately sensitive
Waterlogging	Sensitive
Soil types	Well drained, favouring deep fertile soils
Slope and aspect	Avoid exposed locations
Shade tolerance	Full sun, shade tolerated when very young

Productivity		
Main products	Food	Fruits
	Wood	High-value timber, high yielding
	Biomass	-
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	Low ○
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Moderately dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Risk of toxicity ●
Risks to farming operations		Root suckers Risk of toxicity to other plants, including potatoes, wheat, plum Liable to windthrow and heartrot, especially >60 years
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Native ●
Value to wildlife		Moderate, greatest value for fruits and seeds ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Moderate ●
	Vertebrates	Moderate to high ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Plum (*Prunus domestica* ssp. *domestica*)

Physical			
Typical systems	Arable	Pasture	Lowland
Max. height	Depends on variety and rootstock		
Canopy cover	Globular	Spread depends on variety and rootstock	
Canopy density	Moderately dense		
Root architecture	Shallow and suckering, depending on rootstock		
Growth rate	Moderate		
Silviculture			
Establishment time	5-10 years to maximum height, fruit yields within 4-5 years, full production 7-9 years		
Establishment requirements	Formative pruning	Shelter	Irrigation
Management requirements	Annual pruning in spring to early summer		
Longevity	Unknown		
Rotation length	25-35 years		
Approach to silviculture	Orchard tree		
Tolerances			
High temperatures	Tolerant		
Low temperatures	Hardy in most of UK		
Wind	Sensitive		
Drought	Tolerant		
Waterlogging	Sensitive		
Soil types	Favours sand or clay, acid to neutral, well-drained soils		
Slope and aspect	South or west facing, sheltered		
Shade tolerance	Full sun		

Productivity		
Main products	Food	Fruits – inc. fresh, dehydrated, canned, processed
	Wood	Speciality timber
	Biomass	Limited woodfuel from pruning
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	High ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Moderately dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate (undersized fruits used for feed) ● Risk of toxicity in other plant parts
Risks to farming operations		Root suckers (can be mitigated through appropriate rootstocks)
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Long-established (Archaeophyte – denizen or cultivated) ●
Value to wildlife		Moderate, greatest value for leaf litter, pollen & nectar, and fruits ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	High ●
	Vertebrates	Moderate ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Bird Cherry (*Prunus padus*)

Physical	
Typical systems	Pasture Lowland to semi-upland
Max. height	15-21m
Canopy cover	Spreading, domed and ovoid (conical when young) 4-8m wide
Canopy density	Light to moderately dense
Root architecture	Shallow and suckering
Growth rate	Moderate to high
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Minimal
Management requirements	Prune in mid-summer if silver leaf problematic
Longevity	60 to 100+ years
Rotation length	No typical rotation
Approach to silviculture	Can be coppiced Can be pollarded
Tolerances	
High temperatures	Tolerant
Low temperatures	Hardy
Wind	Conflicting information
Drought	Moderately sensitive
Waterlogging	Moderately tolerant
Soil types	Wide range, favours well-drained damp soils
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	Fruits – inc. for jams, liquer
	Wood	Speciality timber
	Biomass	-
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Low ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Light to moderately dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Risk of toxicity ●
Risks to farming operations		Root suckers
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Native ●
Value to wildlife		Moderate, greatest value for fruits and seeds ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Moderate (high future risk) ●
	Invertebrates	Moderate ●
	Vertebrates	Low to moderate ●
Climate resilience		Low ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Pear (*Pyrus communis*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Depends on variety and rootstock
Canopy cover	Upright and slender to ovoid Spread depends on variety and rootstock
Canopy density	Dense
Root architecture	Generally deep, depending on variety and rootstock
Growth rate	Low to moderate
Silviculture	
Establishment time	Full cropping in 3-9 years (depending on cultivar and rootstock)
Establishment requirements	Staking Formative pruning Water during drought
Management requirements	Annual pruning recommended
Longevity	Potentially 200 to 300 years, depends on cultivar and rootstock
Rotation length	15-40 years , depending on cultivar and rootstock
Approach to silviculture	Orchard tree
Tolerances	
High temperatures	Tolerant
Low temperatures	Hardy
Wind	Sensitive
Drought	Moderately sensitive to moderately tolerant
Waterlogging	Sensitive
Soil types	Wide range, favours well-drained with at least moderate fertility, avoid very acid soils
Slope and aspect	South or west facing, sheltered
Shade tolerance	Full sun, especially when young

Productivity		
Main products	Food	Fruits (inc. fresh, canned, dried, juiced)
	Wood	Speciality timber
	Biomass	Limited woodfuel from pruning
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Moderate ○
	Acidification	Unknown
Shade cover impacts*	Size of shadow (full-grown)	Low, but depends on rootstock/variety ●
	Canopy density	Dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate ○
Risks to farming operations		None known
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Long-established (Archaeophyte – cultivated) ●
Value to wildlife		High, especially for foliage invertebrates, leaf litter, pollen & nectar ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	High ●
	Vertebrates	Moderate ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Sessile Oak (*Quercus petraea*)

Physical			
Typical systems	Arable	Pasture	Lowland to upland
Max. height	Typically up to 27m , exceptionally 40m		
Canopy cover	Globular to broad ovoid, domed	>8m wide, potentially very wide	
Canopy density	Moderately dense		
Root architecture	Deep		
Growth rate	Low to moderate		
Silviculture			
Establishment time	> 50 years to maximum height		
Establishment requirements	Weed control	Protection from browsing	Formative pruning
Management requirements	High pruning for good quality timber		
Longevity	Typically 400-500 years , potentially 600 to 1000+ years		
Rotation length	15-25 years coppice, 120-160 years sawn-wood or veneer		
Approach to silviculture	Can be coppiced (esp. in south)	Can be pollarded	
Tolerances			
High temperatures	Tolerant		
Low temperatures	Very hardy, but occasionally damaged by late frosts		
Wind	Tolerant		
Drought	Moderately tolerant		
Waterlogging	Sensitive		
Soil types	Wide range, favours well-drained somewhat acid soils		
Slope and aspect	Any		
Shade tolerance	Full sun to partial shade		

Productivity		
Main products	Food	-
	Wood	High value timber (inc. sawnwood, veneers, plywood), relatively low yielding
	Biomass	Woodfuel
	Speciality	Acorn flour, various medicinal products
Impact on local soil quality	Nutrient and organic matter accumulation	High ○
	Acidification	High ●
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Moderately dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		Moderate (moderate crude protein, low minerals) ● Risk of toxicity from buds, young leaves and acorns
Risks to farming operations		Capable of becoming very large tree Timber very prone to defects
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Moderate
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Low
Native status		Native ●
Value to wildlife		Very high, especially for mycorrhizal fungi, wood-decay fungi and invertebrates, foliage invertebrates, seeds, and epiphytes ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Moderate ●
	Vertebrates	High ●
Climate resilience		Low to moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Pedunculate Oak (*Quercus robur*)

Physical			
Typical systems	Arable	Pasture	Lowland
Max. height	Typically up to 27m , exceptionally 40m		
Canopy cover	Globular to broad ovoid, domed	>8m wide, potentially very wide	
Canopy density	Moderate to moderately dense		
Root architecture	Deep to very deep, more lateral roots on shallow soils		
Growth rate	Low		
Silviculture			
Establishment time	20-50 years to maximum height		
Establishment requirements	Weed control	Protection from browsing	Formative pruning
Management requirements	High pruning for good quality timber		
Longevity	Typically 400-500 years , potentially 600 to 1000+ years		
Rotation length	15-25 years coppice, 120 or more years sawn-wood or veneer		
Approach to silviculture	Can be coppiced (esp. in south)	Can be pollarded	
Tolerances			
High temperatures	Moderately tolerant		
Low temperatures	Hardy, but occasionally damaged by late frosts		
Wind	Tolerant when established		
Drought	Moderately sensitive to sensitive		
Waterlogging	Moderately sensitive to moderately tolerant		
Soil types	Wide range, favours well-drained but heavy, somewhat base-rich soils		
Slope and aspect	Any		
Shade tolerance	Full sun (especially when young) to partial shade		

Productivity		
Main products	Food	-
	Wood	High value timber (inc. sawnwood, veneers, plywood), relatively low yielding
	Biomass	Woodfuel
	Speciality	Acorn flour, various medicinal products
Impact on local soil quality	Nutrient and organic matter accumulation	High (planted for soil improvement and land restoration) ●
	Acidification	High ●
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Moderate to moderately dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		Moderate (moderate crude protein, low minerals) ● Acorns historically important for pig forage Risk of toxicity from buds, young leaves and acorns
Risks to farming operations		Capable of becoming very large tree Timber very prone to defects
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Moderate
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Low
Native status		Native ●
Value to wildlife		Very high, especially for mycorrhizal fungi, wood-decay fungi and invertebrates, foliage invertebrates, seeds, and epiphytes ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Moderate ●
	Vertebrates	High ●
Climate resilience		Low to moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Red Oak (*Quercus rubra*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	Typically up to 20m , exceptionally 35m
Canopy cover	Globular to broad ovoid >8m wide, potentially >10m
Canopy density	Moderately dense
Root architecture	Likely deep
Growth rate	Moderate to high
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Minimal
Management requirements	High pruning
Longevity	Typically 100 years , potentially 200 years
Rotation length	Typically 70-120 years
Approach to silviculture	Coppices well
Tolerances	
High temperatures	Tolerant
Low temperatures	Very hardy, but occasionally damaged by late frosts
Wind	Tolerant
Drought	Moderately sensitive to tolerant; drought likely to damage timber
Waterlogging	Sensitive
Soil types	Well-drained chalk, sand or clay, favours acid sandy loams
Slope and aspect	Avoid north facing
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	-
	Wood	Timber, lower value than native oaks, relatively low yielding
	Biomass	Woodfuel
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	High ○
	Acidification	High ○
Shade cover impacts*	Size of shadow (full-grown)	Moderate ●
	Canopy density	Moderately dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		Low ○ Risk of toxicity from buds, young leaves and acorns
Risks to farming operations		Capable of becoming very large tree Timber very prone to defects
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Relatively low
	Long-term (60 yrs)	Relatively low
Native status		Recent introduction (Neophyte – naturalised) ●
Value to wildlife		Moderate to high ○
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Moderate ●
	Invertebrates	Low to moderate ●
	Vertebrates	High ○
Climate resilience		High ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

White Willow (*Salix alba*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	25-33m
Canopy cover	Irregular, broadly columnar >8m wide
Canopy density	Open
Root architecture	Extensive, shallow to deep, shallower in wet soils
Growth rate	High
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Protection from browsing Weed control
Management requirements	Strict pruning requirements for timber
Longevity	Typically 20-30 years, exceptionally 200+ years
Rotation length	Timber rotation of 12-20 years for subspecies <i>caerulea</i>
Approach to silviculture	Can be coppiced Can be pollarded
Tolerances	
High temperatures	Moderately tolerant
Low temperatures	Hardy
Wind	Tolerant
Drought	Sensitive
Waterlogging	Tolerant
Soil types	Wide range of damp soils, favouring fertile and base-rich
Slope and aspect	Any
Shade tolerance	Full sun

Productivity		
Main products	Food	-
	Wood	Timber (usually subspecies <i>caerulea</i>), inc. plywood, pulpwood, posts; relatively low yielding
	Biomass	Fuelwood
	Speciality	Tannins, edible truffles
Impact on local soil quality	Nutrient and organic matter accumulation	Low to moderate ○
	Acidification	Low to moderate ○
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate to high (high protein content, moderate digestibility) ●
Risks to farming operations		Risk of blocking drains Winter host of carrot aphid
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Moderate
	Medium-term (40 yrs)	Moderate
	Long-term (60 yrs)	Relatively low
Native status		Long-established (Archaeophyte – denizen) ●
Value to wildlife		Moderate, greatest value for pollen & nectar ●
Other environmental impacts		High water consumption in wet conditions High potential to reduce nutrient leaching
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	High ●
	Vertebrates	Moderate to high ●
Climate resilience		Low ○

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Goat Willow (*Salix caprea*)

Physical	
Typical systems	Arable Pasture Lowland or upland
Max. height	Typically up to 10m , exceptionally 20m
Canopy cover	Irregular, bushy 4-8m wide
Canopy density	Open
Root architecture	Extensive, moderate depth
Growth rate	Moderate to high
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Protection from browsing Weed control
Management requirements	Minimal to no pruning
Longevity	60 to 100+ years
Rotation length	5-12 years
Approach to silviculture	Coppices well
Tolerances	
High temperatures	Moderately tolerant
Low temperatures	Hardy
Wind	Tolerant
Drought	Moderately sensitive
Waterlogging	Moderately sensitive
Soil types	Deep, moist, well-drained
Slope and aspect	Any
Shade tolerance	Full sun to partial shade

Productivity		
Main products	Food	-
	Wood	Speciality timber
	Biomass	Fuelwood, charcoal
	Speciality	Tannins and dye from bark
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Low to moderate ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate to high (high protein content, moderate digestibility) ●
Risks to farming operations		Risk of blocking drains
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Native ●
Value to wildlife		High, especially for foliage invertebrates, pollen & nectar ●
Other environmental impacts		High water consumption in wet conditions High potential to reduce nutrient leaching
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	High ●
	Vertebrates	Moderate to high ●
Climate resilience		Moderate to high ○

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Grey Willow (*Salix cinerea*)

Physical	
Typical systems	Arable Pasture Lowland or upland
Max. height	Typically up to 8m , exceptionally 17m
Canopy cover	Irregular 2.5-4m wide
Canopy density	Open
Root architecture	Extensive, moderate depth
Growth rate	High
Silviculture	
Establishment time	5-10 years to maximum height
Establishment requirements	Protection from browsing Weed control
Management requirements	Minimal to no pruning
Longevity	Up to 100+ years
Rotation length	12-15 years, or short rotation of 2-3 years
Approach to silviculture	Can be coppiced
Tolerances	
High temperatures	Moderately tolerant
Low temperatures	Very hardy
Wind	Tolerant
Drought	Sensitive
Waterlogging	Tolerant
Soil types	Wet and damp soils, favours well-drained chalk, sand or clay
Slope and aspect	Avoid north facing
Shade tolerance	Full sun, more shade tolerant in waterlogged sites

Productivity		
Main products	Food	-
	Wood	-
	Biomass	Fuelwood
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Low to moderate ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate to high ○
Risks to farming operations		Risk of blocking drains
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Native ●
Value to wildlife		High, especially for foliage invertebrates, pollen & nectar ●
Other environmental impacts		High water consumption in wet conditions High potential to reduce nutrient leaching
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	High ●
	Vertebrates	Moderate to high ●
Climate resilience		Low ○

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Willow varieties for SRC (*Salix* spp.)

Physical	
Typical systems	Arable Pasture Lowland or upland (depends on variety)
Max. height	Depends on variety
Canopy cover	Irregular, bushy <4m wide
Canopy density	Open
Root architecture	Extensive, moderate depth
Growth rate	High
Silviculture	
Establishment time	First-year growth typically coppiced
Establishment requirements	Protection from browsing Weed control
Management requirements	Good yields require high inputs of fertiliser, herbicides and pesticides
Longevity	22 to 30 years
Rotation length	2-5 years, typically 3 years
Approach to silviculture	Short-rotation coppice
Tolerances	
High temperatures	Tolerant
Low temperatures	Hardy
Wind	Potentially sensitive
Drought	Sensitive
Waterlogging	Sensitive to frequent waterlogging
Soil types	Wide range, but avoid free-draining sandy soils and high organic peaty soils (due to weed competition)
Slope and aspect	Any
Shade tolerance	Full sun

Productivity		
Main products	Food	-
	Wood	-
	Biomass	Bioenergy, high yielding
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Low to moderate ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Open ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate to high ○
Risks to farming operations		Risk of blocking drains
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	N/A
	Long-term (60 yrs)	N/A
Native status		Hybrids, typically derived from non-native species ●
Value to wildlife		Moderate, greatest value for pollen & nectar ○
Other environmental impacts		High water consumption in wet conditions High potential to reduce nutrient leaching
Resilience		
Pest/disease susceptibility	Diseases	High (but depends on variety) ●
	Invertebrates	High (but depends on variety) ●
	Vertebrates	Moderate to high ●
Climate resilience		Moderate ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Rowan (*Sorbus aucuparia*)

Physical	
Typical systems	Arable Pasture Lowland or upland
Max. height	Typically up to 15m , exceptionally 22m
Canopy cover	Ovoid to domed 4-8m wide
Canopy density	Open to moderately dense
Root architecture	Branching
Growth rate	Moderate
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Protection from browsing Weed control
Management requirements	Minimal to no pruning
Longevity	Up to 200+ years
Rotation length	Not typically grown on rotation
Approach to silviculture	Coppices well
Tolerances	
High temperatures	Moderately sensitive
Low temperatures	Hardy to very hardy
Wind	Tolerant
Drought	Moderately sensitive
Waterlogging	Sensitive
Soil types	Wide range, favours moderately fertile, humus-rich uncompacted soil
Slope and aspect	Any
Shade tolerance	Full sun to partial shade, more shade tolerant when young

Productivity		
Main products	Food	Berries (inc. juicing, jelly, jams)
	Wood	Turnery and speciality timber, low yielding
	Biomass	Limited fuelwood
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	Low ○
	Acidification	Low ○
Shade cover impacts*	Size of shadow (full-grown)	Low ●
	Canopy density	Open to moderately dense ●
	Leaf emergence	Early ●
Livestock fodder benefits		Moderate to high (low crude protein, high digestibility) ●
Risks to farming operations		Winter host for cereal aphid
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	Relatively low
	Medium-term (40 yrs)	Very low
	Long-term (60 yrs)	Very low
Native status		Native ●
Value to wildlife		Moderate, greatest value for leaf litter, blossom, seeds ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	High ●
	Invertebrates	Moderate to high ●
	Vertebrates	High (especially deer) ●
Climate resilience		Low ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

Small-leaved Lime (*Tilia cordata*)

Physical	
Typical systems	Arable Pasture Lowland
Max. height	20-37m
Canopy cover	Broad ovoid to globular, more conical when young >8m wide, potentially >15m
Canopy density	Moderately dense to dense
Root architecture	Conflicting information
Growth rate	Moderate to high
Silviculture	
Establishment time	20-50 years to maximum height
Establishment requirements	Protection from browsing and rodents
Management requirements	Minimal to no pruning
Longevity	800+ years, potentially exceeding 1000 years if coppiced
Rotation length	10-20 years coppice, 100-140 years timber
Approach to silviculture	Coppices well Can be pollarded
Tolerances	
High temperatures	Moderately tolerant
Low temperatures	Hardy to very hardy
Wind	Sensitive
Drought	Moderately sensitive, considerably reduces growth
Waterlogging	Sensitive
Soil types	Wide range, favours neutral to slightly alkaline, moist and fertile soils
Slope and aspect	Favours valley bottoms with moist air, avoid exposed locations
Shade tolerance	Full sun to partial shade or greater

Productivity		
Main products	Food	-
	Wood	Timber, moderate yielding
	Biomass	-
	Speciality	-
Impact on local soil quality	Nutrient and organic matter accumulation	High (often planted as soil improver) ●
	Acidification	Low ●
Shade cover impacts*	Size of shadow (full-grown)	High ●
	Canopy density	Moderately dense to dense ●
	Leaf emergence	Late ●
Livestock fodder benefits		High (high nutritional value and digestibility) ●
Risks to farming operations		None known
Environmental impacts		
Carbon sequestration	Short-term (20 yrs)	High
	Medium-term (40 yrs)	High
	Long-term (60 yrs)	High
Native status		Native ●
Value to wildlife		Moderate, greatest value for mycorrhizal fungi, leaf litter, pollen & nectar ●
Other environmental impacts		None known
Resilience		
Pest/disease susceptibility	Diseases	Low to moderate ●
	Invertebrates	Moderate ●
	Vertebrates	High (especially browsing mammals) ●
Climate resilience		High ●

* Effect depends on system, likely benefits of shade in livestock systems but disbenefit in arable
● = high confidence, ● = moderate confidence, ○ = low confidence

References

- Alexander, K., Butler, J., Green, T. (2006) The value of different tree and shrub species to wildlife. *British Wildlife*. **18**(1), 18–28.
- Andrianarisoa, K.S., Dufour, L., Bienaimé, S., Zeller, B., Dupraz, C. (2016) The introduction of hybrid walnut trees (*Juglans nigra* × *regia* cv. NG23) into cropland reduces soil mineral N content in autumn in southern France. *Agroforestry Systems*. **90**(2), 193–205.
- Bentrup, G., Hopwood, J., Adamson, N.L., Vaughan, M. (2019) Temperate agroforestry systems and insect pollinators: A review. *Forests*. **10**(981), 1–20.
- Böhm, C., Kanzler, M., Freese, D. (2014) Wind speed reductions as influenced by woody hedgerows grown for biomass in short rotation alley cropping systems in Germany. *Agroforestry Systems*. **88**(4), 579–591.
- Briggs, S., Knight, I. (2019) The economic case for agroforestry. In B. Raskin & S. Osborn, eds. *The Agroforestry Handbook*. Bristol: Soil Association Limited, pp. 95–141.
- Casals, P., Romero, J., Rusch, G.M., Ibrahim, M. (2014) Soil organic C and nutrient contents under trees with different functional characteristics in seasonally dry tropical silvopastures. *Plant and Soil*. **374**(1–2), 643–659.
- Castro-Díez, P., Vaz, A.S., Silva, J.S., van Loo, M., Alonso, Á., Aponte, C., Bayón, Á., Bellingham, P.J., Chiuffo, M.C., DiManno, N., Julian, K., Kandert, S., La Porta, N., Marchante, H., Maule, H.G., Mayfield, M.M., Metcalfe, D., Monteverdi, M.C., Núñez, M.A., Ostertag, R., Parker, I.M., Peltzer, D.A., Potgieter, L.J., Raymundo, M., Rayome, D., Reisman-Berman, O., Richardson, D.M., Roos, R.E., Saldaña, A., Shackleton, R.T., Torres, A., Trudgen, M., Urban, J., Vicente, J.R., Vilà, M., Ylioja, T., Zenni, R.D., Godoy, O. (2019) Global effects of non-native tree species on multiple ecosystem services. *Biological Reviews*. **94**(4), 1477–1501.
- CEH Ammonia reduction calculator. <https://farmtreestoair.ceh.ac.uk/ammonia-reduction-calculator>.
- Hagen-Thorn, A., Callesen, I., Armolaitis, K., Nihlgård, B. (2004) The impact of six European tree species on the chemistry of mineral topsoil in forest plantations on former agricultural land. *Forest Ecology and Management*. **195**(3), 373–384.
- Hall, R., Allen, S., Rosier, P., Smith, D., Hodnett, M., Roberts, J., Hopkins, R., Davies, H. (1996) *Hydrological Effects of Short Rotation Energy Coppice*. Wallingford: Institute of Hydrology and British Geological Survey.
- Herbst, M., Eschenbach, C., Kappen, L. (1999) Water use in neighbouring stands of beech (*Fagus sylvatica* L.) and black alder (*Alnus glutinosa* (L.) Gaertn.). *Annals of Forest Science*. **56**(2), 107–120.
- Hill, A.R. (2019) Groundwater nitrate removal in riparian buffer zones: a review of research progress in the past 20 years. *Biogeochemistry*. **143**(3), 347–369.

- Hotchkiss, A., Herbert, S. (2022) *Tree species handbook*. Grantham, UK: The Woodland Trust.
- Isaac, M.E., Borden, K.A. (2019) Nutrient acquisition strategies in agroforestry systems. *Plant and Soil*. **444**(1–2), 1–19.
- Jug, A., Makeschin, F., Rehfuess, K.E., Hofmann-Schielle, C. (1999) Short-rotation plantations of balsam poplars, aspen and willows on former arable land in the Federal Republic of Germany. III. Soil ecological effects. *Forest Ecology and Management*. **121**(1–2), 85–99.
- Kassa, G., Bekele, T., Demissew, S., Abebe, T. (2022) Leaves litterfall and nutrient inputs from four multipurpose tree/shrub species of homegarden agroforestry systems. *Environmental Systems Research* 2022 11:1. **11**(1), 1–16.
- Kendall, N.R., Smith, J., Whistance, L.K., Stergiadis, S., Stoate, C., Chesshire, H., Smith, A.R. (2021) Trace element composition of tree fodder and potential nutritional use for livestock. *Livestock Science*. **250**, 104560.
- Kim, D.G., Isaac, M.E. (2022) Nitrogen dynamics in agroforestry systems. A review. *Agronomy for Sustainable Development* 2022 42:4. **42**(4), 1–18.
- Kletty, F., Rozan, A., Hibold, C. (2023) Biodiversity in temperate silvoarable systems: A systematic review. *Agriculture, Ecosystems & Environment*. **351**, 108480.
- Laganière, J., Angers, D.A., Paré, D. (2010) Carbon accumulation in agricultural soils after afforestation: A meta-analysis. *Global Change Biology*. **16**(1), 439–453.
- Mayer, P.M., Reynolds, S.K., McCutchen, M.D., Canfield, T.J. (2007) Meta-Analysis of Nitrogen Removal in Riparian Buffers. *Journal of Environmental Quality*. **36**(4), 1172–1180.
- McAdam, J., Olave, R., Fornara, D. (2018) Silvopastoral agroforestry - an option to support sustainable grassland intensification. In *4th European Agroforestry Conference*. pp. 178–180.
- Mmolotsi, R. (2004) *Potential uses of red alder (Alnus rubra Bong) in silvopastoral systems*. Bangor: Bangor University.
- Nisbet, T. (2005) *Water Use by Trees*. Edinburgh: Forestry Commission.
- Nisbet, T., Evans, C. (2014) *Forestry and Surface Water Acidification*. Farnham: Forest Research.
- Nisbet, T., Silgram, M., Shah, N., Morrow, K., Broadmeadow, S. (2011) *Woodland for water: woodland measures for meeting water framework directive objectives*. Forest Research.
- Ola, A., Dodd, I.C., Quinton, J.N. (2015) Can we manipulate root system architecture to control soil erosion? *Soil*. **1**(2), 603–612.
- Pardon, P., Reubens, B., Reheul, D., Mertens, J., De Frenne, P., Coussement, T., Janssens, P., Verheyen, K. (2017) Trees increase soil organic carbon and nutrient

availability in temperate agroforestry systems. *Agriculture, Ecosystems & Environment*. **247**, 98–111.

Purahong, W., Kapturska, D., Pecyna, M.J., Schulz, E., Schloter, M., Buscot, F., Hofrichter, M., Kružer, D. (2014) Influence of Different Forest System Management Practices on Leaf Litter Decomposition Rates, Nutrient Dynamics and the Activity of Ligninolytic Enzymes: A Case Study from Central European Forests. *PLOS ONE*. **9**(4), e93700.

Raskin, B., Osborn, S. eds. (2019) *The Agroforestry Handbook*. Bristol: Soil Association.

Regni, L., Bartucca, M.L., Pannacci, E., Tei, F., Del Buono, D., Proietti, P. (2021) Phytodepuration of Nitrate Contaminated Water Using Four Different Tree Species. *Plants* **2021**, Vol. 10, Page 515. **10**(3), 515.

Reubens, B., Poesen, J., Danjon, F., Geudens, G., Muys, B. (2007) The role of fine and coarse roots in shallow slope stability and soil erosion control with a focus on root system architecture: A review. *Trees - Structure and Function*. **21**(4), 385–402.

Savill, P. (2019) *The Silviculture of Trees Used in British Forestry*. 3rd edition. Wallingford: CABI.

De Schrijver, A., De Frenne, P., Staelens, J., Verstraeten, G., Muys, B., Vesterdal, L., Wuyts, K., van Nevel, L., Schelfhout, S., De Neve, S., Verheyen, K. (2012) Tree species traits cause divergence in soil acidification during four decades of postagricultural forest development. *Global Change Biology*. **18**(3), 1127–1140.

Sjoman, H., Anderson, A. (2023) *The Essential Tree Selection Guide*. Filbert Press and Royal Botanic Gardens, Kew.

Soil Association (2023) *Investigating the feasibility for an Agroforestry Carbon Code - NEIRF Phase 2 - Final report and recommendations*.

Sollen-Norrlin, M., Ghaley, B.B., Rintoul, N.L.J. (2020) Agroforestry benefits and challenges for adoption in Europe and beyond. *Sustainability*. **12**(7001), 1–20.

Stace, C.A. (2019) *New Flora of the British Isles*. 4th ed. Suffolk: C & M Floristics.

Stokes, A., Douglas, G.B., Fourcaud, T., Giadrossich, F., Gillies, C., Hubble, T., Kim, J.H., Loades, K.W., Mao, Z., Mclvor, I.R., Mickovski, S.B., Mitchell, S., Osman, N., Phillips, C., Poesen, J., Polster, D., Preti, F., Raymond, P., Rey, F., Schwarz, M., Walker, L.R. (2014) Ecological mitigation of hillslope instability: Ten key issues facing researchers and practitioners. *Plant and Soil*. **377**(1–2), 1–23.

Tang, Y.S., Mansfield, P., Braban, C.F., Brass, D., Davies, R., Arkle, P., Stephens, A., Iwanicka, A., Mullinger, N.J., Leeson, S., Cowan, N., Simmons, I., Martin Hernandez, C., Jones, R., Fisher, B., Shutt, M., Timmis, R., Whyatt, D., Vanguelova, E., Benham, S., Broadmeadow, S., Bealey, W.J. (2022) *Ammonia Reduction by Trees: Field case studies for monitoring ammonia reduction by treebelts*. Penicuik: UK Centre for Ecology and Hydrology.

Teklehaimanot, Z., Jones, M., Sinclair, F.L. (2002) Tree and livestock productivity in relation to tree planting configuration in a silvopastoral system in North Wales, UK. *Agroforestry Systems*. **56**(1), 47–55.

Torralba, M., Fagerholm, N., Burgess, P.J., Moreno, G., Plieninger, T. (2016) Do European agroforestry systems enhance biodiversity and ecosystem services? A meta-analysis. *Agriculture, Ecosystems & Environment*. **230**, 150–161.

Ucar, T., Hall, F.R. (2001) Windbreaks as a pesticide drift mitigation strategy: a review. *Pest Management Science*. **57**(8), 663–675.

Ucar, T., Hall, F.R., Tew, J.E., Hacker, J.K. (2003) Wind tunnel studies on spray deposition on leaves of tree species used for windbreaks and exposure of honey bees. *Pest Management Science*. **59**(3), 358–364.

Vannoppen, W., De Baets, S., Keeble, J., Dong, Y., Poesen, J. (2017) How do root and soil characteristics affect the erosion-reducing potential of plant species? *Ecological Engineering*. **109**, 186–195.

Wenneker, M., Van de Zande, J. (2008) Spray drift reducing effects of natural windbreaks in orchard spraying. *Aspects of Applied Biology*. **84**, 1–8.

Authors

Tom Staton (University of Reading), Kate Beauchamp (Forest Research), Alice Broome (Forest Research), Tom Breeze (University of Reading).

Acknowledgements

A draft report, or parts of it, were reviewed by: Rory Lunny (Defra), Daniel Hewitt and Vanessa Burton (The Woodland Trust), Geoff Newman (Natural England), Tom Nisbet, Suzanne Robinson and Elena Vanguelova (Forest Research). We are grateful to the stakeholder group who provided valuable input throughout the development of this guide.